

IVIG Intake Form (Autoimmune)



Demographic Information

Last Name:		First Name:		SS#:	
Home Address:			City:	State:	Zip:
DOB:	Height:	Weight:	Allergies:		
Phone (H):		Phone (C):		Phone (W):	
Emergency Contact: _____			Patient's Gender: <input type="checkbox"/> Male <input type="checkbox"/> Female		
Phone (H): _____		Phone (C): _____		Phone (W): _____	

Insurance Information

Primary Insurance:		Subscriber Name:	
Policy #:	Group #:	PCN:	
RX BIN:	Insurance Phone:		
Secondary Insurance:		Subscriber Name:	
Policy #:	Group #:	PCN:	
RX BIN:	Insurance Phone:		

Diagnosis

<input type="checkbox"/> 357.81 Chronic Inflammatory Demyelinating Polyneuropathy	<input type="checkbox"/> 357.0 Acute Infective Polyneuritis (Guillan-Barre Syndrome)	<input type="checkbox"/> 356.4 Polyneuropathy Idiopathic, Progressive
<input type="checkbox"/> 287.31 ITP	<input type="checkbox"/> 358.0 Myasthenia Gravis	<input type="checkbox"/> 357.9 Multifocal Motor Neuropathy
<input type="checkbox"/> 357.82 Critical Illness Polyneuropathy (acute motor neuropathy)	<input type="checkbox"/> 710.0 Systemic Lupus Erythematosus	<input type="checkbox"/> 694.4 Pemphigus Vulgaris
<input type="checkbox"/> 710.3 Dermatomyositis	<input type="checkbox"/> 710.4 Polymyositis	<input type="checkbox"/> 694.4 Pemphigus Foliaceus
<input type="checkbox"/> 333.91 Stiff-man Syndrome	<input type="checkbox"/> 357.2 Acute Infective Polyneuritis	<input type="checkbox"/> 694.5 Bullous Pemphigoid
<input type="checkbox"/> 340.0 Multiple Sclerosis (Relapsing / Remitting)	<input type="checkbox"/> 446.1 Kawasaki Syndrome	<input type="checkbox"/> 696.0 Psoriatic Arthropathy
<input type="checkbox"/> Other: _____	<input type="checkbox"/> ICD9: _____	

Additional Medical History

History of: ☐ Renal Insufficiency ☐ Diabetes ☐ HTN ☐ CHF ☐ Thromboembolic Event
☐ Other: _____

Physician Information

MD:	License:	NPI:
Hospital/Clinic:		Phone:
Fax:	Address:	
Office Contact:		Medicaid #:

Referral Phone: 787-704-2025

Referral Fax: 787-704-2027

Patient Name: _____ Date of Birth: _____

Prescription & Product Information

■ May Not Substitute

IVIG Product:	Concentration %:
Dose: _____ mg/kg (+/- 10%) = _____ grams	
Frequency: Repeat dose <input type="checkbox"/> Daily for _____ days every _____ weeks <input type="checkbox"/> Weekly <input type="checkbox"/> Monthly Other: _____ x _____ doses # of Refills _____	
Administration Rate: <input type="checkbox"/> Per manufacturer guidelines, as tolerated <input type="checkbox"/> Other _____	
<i>*Patients at risk for renal involvement will be run at a conservative rate*</i>	
Has the patient received IVIG previously? <input type="checkbox"/> Yes <input type="checkbox"/> No	Date of last dose:

Pre Medications

<input type="checkbox"/> Hydration: <input type="checkbox"/> Prior to <input type="checkbox"/> During <input type="checkbox"/> Following infuse: _____ mL _____ solution
<input type="checkbox"/> Benadryl: <input type="checkbox"/> 25mg <input type="checkbox"/> 50mg 15-25 minutes prior to infusion <input type="checkbox"/> PO <input type="checkbox"/> IVP
<input type="checkbox"/> Tylenol: <input type="checkbox"/> 650mg <input type="checkbox"/> 1000mg 15-25 minutes prior to infusion PO
<input type="checkbox"/> Solu-Cortef 100mg/2mL <input type="checkbox"/> Solu-Medrol 125mg/2mL _____ mg slow IVP: <input type="checkbox"/> pre <input type="checkbox"/> halfway <input type="checkbox"/> upon completion
<input type="checkbox"/> Lidocaine cream 4% applied topically for venous or port access
Other:

Delivery Method

<input type="checkbox"/> Gravity using rate control device <input type="checkbox"/> Pump	Vascular Device: <input type="checkbox"/> PIV <input type="checkbox"/> Central: _____
--	---

Flushes

Normal Saline: <input type="checkbox"/> 5mL <input type="checkbox"/> 10mL <input type="checkbox"/> Pre/Post Infusion <input type="checkbox"/> 20mL <input type="checkbox"/> Post infusion
Heparin: _____ mL (_____ u/mL) as SASH

Nursing Assessment

- ☐ -Skilled nursing visit to: establish IV access, medication administration as prescribed, provide patient education related to disease state/therapy, assess general status and response to therapy. Frequency determined by therapy schedule.
- ☐ -Obtain baseline vital signs
- ☐ -Monitor vital signs every 15 minutes for first hour. Every 30 minutes for second hour. Then hourly to completion.
- ☐ -Provide needles, syringes, VAD and other ancillary supplies required for safe infusion.
- ☐ -Discontinue use and notify prescribing physician if patient demonstrates any of the following:
Fluid overload, cardiovascular symptoms, allergic reaction, moderate/severe headache, s/sx Aseptic Meningitis.
- ☐ -RN to draw following labs: _____
Frequency _____

Procedure for Anaphylaxis (pharmacy to provide)

1. Stop Infusion
2. Call 911 and prescribing physician immediately
3. Administer the following: ☐ Diphenhydramine 25-50mg slow IVP Q 4 hours PRN, dispense (1) 50mg vial
☐ Diphenhydramine 25mg capsules, use as directed, dispense 4 capsules
☐ Epinephrine (1:1000) 0.4mg subcutaneously PRN for anaphylaxis, dispense 1
☐ Normal Saline 500ml, use as directed, dispense 1 bag

I authorize SPS Specialty Pharmacy Services and its representatives to act as an agent to initiate and execute the insurance prior authorization process.

Physician Signature: _____ Date: _____

Referral Phone: 787-704-2025

Referral Fax: 787-704-2027